


1. An antimycobacterial compound which comprises the formula:



5 wherein  $R_1$  is H; and

$R_2$  is  $C_3$  to  $C_{14}$  alkyl,  $C_3$  to  $C_{10}$  substituted alkyl,  $C_2$  to  $C_{10}$  alkenyl,  $C_2$  to  $C_9$  substituted alkenyl,  $C_2$  to  $C_9$  substituted dialkenyl,  $C_3$  to  $C_7$  cycloalkyl,  $C_3$  to  $C_7$  substituted cycloalkyl, phenyl, substituted phenyl,  $C_7$  to  $C_{16}$  phenylalkyl,  $C_7$  to  $C_{16}$  substituted phenylalkyl, benzyl, substituted benzyl, naphthyl, substituted naphthyl, heterocycle, substituted heterocycle, halo, hydroxy, amino, or carboxy;

or a pharmaceutically acceptable salt thereof; or a pharmaceutical isomer thereof; or a combination of the same.

2. The antimycobacterial compound according to claim 1 wherein  $R_1$  is H; and

5  $R_2$  is  $CH=CHCH_3$  (trans),  $CH=CHCH_2CH_3$  (trans),  $CH=CHCH_2CH_2CH_3$  (trans),  $CH=CHCH_2CH_2CH_2CH_3$  (trans),  $C(CH_3)=CHCH_3$  (trans),  $CH=C(CH_3)CH_2CH_2CH=C(CH_3)_2$  (trans),  $CH=NNHCO-4-C_6H_4NHCH_2CH(CH_3)CH_2CH_2CH=C(CH_3)_2$ ,  $4-C_6H_4-CH=NNHCO-4-C_5H_4N$ ,  $4-C_6H_4-O-CH_2CH_2CH_2CH_3$ ,  $(CH_2)_{11}CH_3$ ,  $4-C_6H_4NO_2$ ,  $C_6H_5$ ,  $2-C_6H_4OH$ ,  $4-OH-3-OCH_3C_6H_3$ ,  $4-C_6H_4OCH_3$ ,  $3-C_6H_4OCH_3$ ,  $(CH_2)_8CH_3$ ,  $(CH_2)_2CH_3$ ,  $2-C_6H_4OCH_3$ ,  
10  $C(CH_3)=CHC_6H_5$  (trans),  $4-C_6H_4F$ ,  $3,5-di(CH_3)-4-O-C_7H_7$ ,  $2-F-4-OCH_3C_6H_3$ ,  $2-ClC_6H_4$ ,  $4-BrC_6H_4$ ,  $3-C_6H_4NO_2$ ,  $4-C_6H_4O(CH_2)_5CH_3$ ,  $2-Cl-5-NO_2C_6H_3$ ,  $4-Cl-3-NO_2C_6H_3$ ,  $2-C_6H_4NO_2$ ,  $2-6-$

[illegible]

all compound acco

all compound acco

all compound acco

all compound acco

all compound acco

all compound acco

all compound acco

all compound acco

all compound acco

all compound acco


54b  
02

7. The ar  
4C<sub>6</sub>H<sub>8</sub>NNHCC

54b  
02



7. The ar  
4C<sub>6</sub>H<sub>8</sub>NNHCC

8. The antimycobacterial compound according to claim 1 where  $R_1, R_2$  is




Sub  
22  
cont

or





5

9. The antimycobacterial compound according to claim 1 wherein  $R_1$  is H; and  $R_2$  is



10

Cc1ccc2cc(OC)ccc2c1

wherein R<sub>1</sub> is H; R<sub>2</sub> is C<sub>3</sub> to C<sub>14</sub> alkyl, C<sub>3</sub> to C<sub>10</sub> substituted alkyl, C<sub>2</sub> to C<sub>10</sub> alkenyl, C<sub>2</sub> to C<sub>9</sub> substituted alkenyl, C<sub>2</sub> to C<sub>9</sub> substituted dialkenyl, C<sub>3</sub> to C<sub>7</sub> cycloalkyl, C<sub>3</sub> to C<sub>7</sub> substituted cycloalkyl, phenyl, substituted phenyl, C<sub>7</sub> to C<sub>16</sub> phenylalkyl, C<sub>7</sub> to C<sub>16</sub> substituted phenylalkyl, benzyl, substituted benzyl, naphthyl, substituted naphthyl, heterocycle, substituted heterocycle; and

R<sub>3</sub> is C<sub>1</sub> or C<sub>2</sub> alkyl; or a pharmaceutically acceptable salt thereof; or a pharmaceutically acceptable isomer thereof; or a combination of the same.

11. The antimycobacterial compound according to claim 10 wherein  $R_1$  is H;  $R_2$  is 2,6-di(Cl) $C_6H_3$ , 3- $NO_2$ -4-Cl- $C_6H_3$ , 3,4-di(F) $C_6H_3$ , 2- $C_6H_4NO_2$ , 3,4-di(Cl) $C_6H_3$  and 2,6-di(F) $C_6H_3$ ;

and

$R_3$  is  $CH_3$ .

12. The antimycobacterial compounds according to claim 10 wherein  $R_1$  is  $CH_3$ ;  $R_2$  is  $CH_3$ ; and

$R_3$  is  $CH_2CH_3$  or  $CH_3$ .

13. The antimycobacterial compounds according to claim 10 wherein  $R_1$ ,  $R_2$  is  $(CH_2)_5$ ;


and

$R_3$  is  $CH_3$ .

14. The antimycobacterial compound according to claim 10 wherein  $R_1$  is  $CH_3$ ;  $R_2$  is  $C_6H_5$ ; and

$R_3$  is  $CH_3$ .

15. An antimycobacterial compound which comprises the formula:




III

- 5 wherein  $R_1$  is  $C_2$  to  $C_6$  alkyl,  $C_2$  to  $C_6$  substituted alkyl,  $C_2$  to  $C_{10}$  alkenyl,  $C_2$  to  $C_{10}$  substituted alkenyl,  $C_3$  to  $C_7$  cycloalkyl,  $C_3$  to  $C_7$  substituted cycloalkyl, phenyl, substituted phenyl,  $C_7$  to  $C_{16}$  phenylalkyl,  $C_7$  to  $C_{16}$  substituted phenylalkyl, benzyl, substituted benzyl, naphthyl, substituted naphthyl, heterocycle, substituted heterocycle;
- 10 or a pharmaceutically acceptable salt thereof; or a pharmaceutically acceptable isomer thereof; or a combination of the same.

16. The antimycobacterial compound according to claim 15 wherein  $R_1$  is  $NHC_6H_5$ ,  $NH-4-C_6H_4CH_3$ ,  $NH-4-C_6H_4Br$  or  $NH-4-C_6H_4Cl$ .

17. A method for producing an antimycobacterial compound comprising the formula of:



- 5 wherein  $R_1$  is H or  $CH_3$ ; and

wherein  $R_2$  is  $C_1$  to  $C_{14}$  alkyl,  $C_2$  to  $C_{10}$  substituted alkyl,  $C_2$  to  $C_{10}$  alkenyl,  $C_2$  to  $C_9$  substituted alkenyl,  $C_2$  to  $C_9$  substituted dialkenyl,  $C_3$  to  $C_7$  cycloalkyl,  $C_3$  to  $C_7$  substituted cycloalkyl, phenyl, substituted phenyl,  $C_7$  to  $C_{16}$  phenylalkyl,  $C_7$  to  $C_{16}$  substituted phenylalkyl, benzyl, substituted benzyl, naphthyl, substituted naphthyl, heterocycle, substituted heterocycle, halo, hydroxy, amino, or carboxy; or


10

wherein  $R_1R_2 = C_4$  to  $C_8$  cycloalkyl or  $C_4$  to  $C_{10}$  substituted cycloalkyl;

which comprises:

15

refluxing



(1)

with absolute ethanol to produce a solution;

adding a carbonyl compound comprising the formula of:

wherein  $R_3 = H$  or  $CH_3$ ; and

wherein  $R_4 = C_1$  to  $C_{14}$  alkyl,  $C_2$  to  $C_{10}$  substituted alkyl,  $C_2$  to  $C_{10}$  alkenyl,  $C_2$  to  $C_9$  substituted alkenyl,  $C_2$  to  $C_9$  substituted dialkenyl,  $C_3$  to  $C_7$  cycloalkyl,  $C_3$  to  $C_7$  substituted cycloalkyl, phenyl, substituted phenyl,  $C_7$  to  $C_{16}$  phenylalkyl,  $C_7$  to  $C_{16}$  substituted phenylalkyl, benzyl, substituted benzyl, naphthyl, substituted naphthyl, heterocycle, substituted heterocycle, halo, hydroxy, amino, or carboxy; or

wherein  $R_3R_4 = C_4$  to  $C_8$  cycloalkyl or  $C_4$  to  $C_{10}$  substituted cycloalkyl;

to the solution to produce a reaction mixture;

distilling the reaction mixture;


adding diethyl ether to the reaction mixture;

filtering the reaction mixture; and

drying the filtrate to produce I.



18. A method for producing an antimycobacterial compound comprising the formula of:



wherein  $R_1$  = wherein  $R_1$  is H or  $CH_3$


wherein  $R_2$  =  $C_1$  to  $C_{14}$  alkyl,  $C_2$  to  $C_{10}$  substituted alkyl,  $C_2$  to  $C_{10}$  alkenyl,  $C_2$  to  $C_9$  substituted alkenyl,  $C_2$  to  $C_9$  substituted dialkenyl,  $C_3$  to  $C_7$  cycloalkyl,  $C_3$  to  $C_7$  substituted cycloalkyl, phenyl, substituted phenyl,  $C_7$  to  $C_{16}$  phenylalkyl,  $C_7$  to  $C_{16}$  substituted phenylalkyl, benzyl, substituted benzyl, naphthyl, substituted naphthyl, heterocycle, substituted heterocycle, halo, hydroxy, amino, or carboxy; or

wherein  $R_1R_2$  =  $C_4$  to  $C_8$  cycloalkyl or  $C_4$  to  $C_{10}$  substituted cycloalkyl;

wherein  $R_3$  =  $C_1$  or  $C_2$  alkyl

which comprises:

refluxing




wherein  $R_1$  is H or  $CH_3$ ; and

wherein  $R_2$  is  $C_1$  to  $C_{14}$  alkyl,  $C_2$  to  $C_{10}$  substituted alkyl,  $C_2$  to  $C_{10}$  alkenyl,  $C_2$  to  $C_9$  substituted alkenyl,  $C_2$  to  $C_9$  substituted dialkenyl,  $C_3$  to  $C_7$  cycloalkyl,  $C_3$  to  $C_7$  substituted cycloalkyl, phenyl, substituted phenyl,  $C_7$  to  $C_{16}$  phenylalkyl,  $C_7$  to  $C_{16}$  substituted phenylalkyl, benzyl, substituted benzyl, naphthyl, substituted naphthyl, heterocycle, substituted heterocycle, halo, hydroxy, amino, or carboxy; or

wherein  $R_1R_2 = C_4$  to  $C_8$  cycloalkyl or  $C_4$  to  $C_{10}$  substituted cycloalkyl;

with a carboxylic acid anhydride comprising the formula of:



wherein  $R = C_1$  or  $C_2$  alkyl

to produce a reaction mixture;

drying the reaction mixture;


adding ether to the reaction mixture to form a solution;

separating the ether from the solution to yield an aqueous layer;

extracting the aqueous layer with ether;

drying the ether extracts to produce II.

19. A method for producing a compound comprising the formula of:




III

wherein  $R_1 = C_1$  to  $C_6$  alkyl,  $C_2$  to  $C_6$  substituted alkyl,  $C_2$  to  $C_{10}$  alkenyl,  $C_2$  to  $C_{10}$  substituted alkenyl,  $C_3$  to  $C_7$  cycloalkyl,  $C_3$  to  $C_7$  substituted cycloalkyl, phenyl, substituted phenyl,  $C_7$  to  $C_{16}$  phenylalkyl,  $C_7$  to  $C_{16}$  substituted phenylalkyl, benzyl, substituted benzyl, naphthyl, substituted naphthyl, heterocycle, substituted heterocycle;

which comprises:


refluxing



(1)

with ethanol to produce a solution;

adding an isothiocyanate comprised of the formula of:



wherein  $R = C_1$  to  $C_6$  alkyl,  $C_2$  to  $C_6$  substituted alkyl,  $C_2$  to  $C_{10}$  alkenyl,  $C_2$  to  $C_{10}$  substituted alkenyl,  $C_3$  to  $C_7$  cycloalkyl,  $C_3$  to  $C_7$  substituted cycloalkyl, phenyl, substituted phenyl,  $C_7$  to  $C_{16}$


phenylalkyl, C<sub>7</sub> to C<sub>16</sub> substituted phenylalkyl, benzyl, substituted benzyl, naphthyl, substituted naphthyl, heterocycle, substituted heterocycle;

to the solution to form a reaction mixture;

cooling the reaction mixture;

filtering the reaction mixture to produce **III**.


20. A method for producing an antimycobacterial compound comprising the formula of:



wherein R<sub>1</sub> = C<sub>1</sub> to C<sub>6</sub> alkyl, C<sub>2</sub> to C<sub>6</sub> substituted alkyl, C<sub>2</sub> to C<sub>10</sub> alkenyl, C<sub>2</sub> to C<sub>10</sub> substituted alkenyl, C<sub>3</sub> to C<sub>7</sub> cycloalkyl, C<sub>3</sub> to C<sub>7</sub> substituted cycloalkyl, phenyl, substituted phenyl, C<sub>7</sub> to C<sub>16</sub> phenylalkyl, C<sub>7</sub> to C<sub>16</sub> substituted phenylalkyl, benzyl, substituted benzyl, naphthyl, substituted naphthyl, heterocycle, substituted heterocycle;

which comprises:

adding diethyl ether to




(I)

to produce a solution;

boiling the solution;

adding a carboxylic acid anhydride comprising the formula of:



wherein R = C<sub>1</sub> to C<sub>6</sub> alkyl, C<sub>2</sub> to C<sub>6</sub> substituted alkyl, C<sub>2</sub> to C<sub>10</sub> alkenyl, C<sub>2</sub> to C<sub>10</sub> substituted alkenyl, C<sub>3</sub> to C<sub>7</sub> cycloalkyl, C<sub>3</sub> to C<sub>7</sub> substituted cycloalkyl, phenyl, substituted phenyl, C<sub>7</sub> to C<sub>16</sub> phenylalkyl, C<sub>7</sub> to C<sub>16</sub> substituted phenylalkyl, benzyl, substituted benzyl, naphthyl, substituted naphthyl, heterocycle, substituted heterocycle;

in ether to the solution to form a reaction mixture;

refluxing the reaction mixture;

cooling the reaction mixture to produce IV.

add  
a4

B2